

337 U.S. PTO

A Boxseq +

Please type a plus sign (+) inside this box → ☐

PTO/SB/05 (2/98)
Approved for use through 09/30/2000. OMB 0651-0032
Patent and Trademark Office: U.S. DEPARTMENT OF COMMERCE
Under the Paperwork Reduction Act of 1995, no persons are required to respond to a collection of information unless it displays a valid OMB control number.

UTILITY PATENT APPLICATION TRANSMITTAL

(Only for new nonprovisional applications under 37 C.F.R. § 1.53(b))

Attorney Docket No.	49321-1
First Inventor or Application Identifier	Doherty, Joni K.
Title	HER-2 BINDING ANTAGONISTS
Express Mail Label No.	EM563437079US

APPLICATION ELEMENTS

See MPEP chapter 600 concerning utility patent application contents.

ADDRESS TO: Assistant Commissioner for Patents
Box Patent Application
Washington, DC 20231

- ☐ * Fee Transmittal Form (e.g., PTO/SB/17)
(Submit an original and a duplicate for fee processing)
- ☒ Specification [Total Pages 33]
(preferred arrangement set forth below)
 - Descriptive title of the invention
 - Cross References to Related Applications
 - Statement Regarding Fed sponsored R & D
 - Reference to Microfiche Appendix
 - Background of the invention
 - Brief Summary of the invention
 - Brief Description of the Drawings (if filed)
 - Detailed Description
 - Claim(s)
 - Abstract of the Disclosure
- ☒ Drawing(s) (35 U.S.C. 113) [Total Sheets 7]
- Oath or Declaration [Total Pages 2]
 - ☐ Newly executed (original or copy)
 - ☐ Copy from a prior application (37 C.F.R. § 1.63(d))
(for continuation/divisional with Box 17 completed)
(Note Box 5 below)
 - ☐ DELETION OF INVENTOR(S)
Signed statement attached deleting inventor(s) named in the prior application, see 37 C.F.R. §§ 1.63(d)(2) and 1.33(b).
- ☐ Incorporation By Reference (useable if Box 4b is checked)
The entire disclosure of the prior application, from which a copy of the oath or declaration is supplied under Box 4b, is considered to be part of the disclosure of the accompanying application and is hereby incorporated by reference therein.

- ☐ Microfiche Computer Program (Appendix)
- Nucleotide and/or Amino Acid Sequence Submission (if applicable, all necessary)
 - ☒ Computer Readable Copy
 - ☒ Paper Copy (identical to computer copy)
 - ☒ Statement verifying identity of above copies

ACCOMPANYING APPLICATION PARTS

- ☒ Assignment Papers (cover sheet & document(s))
- ☐ 37 C.F.R. § 3.73(b) Statement (when there is an assignee) ☒ Power of Attorney
- ☐ English Translation Document (if applicable)
- ☐ Information Disclosure Statement (IDS)/PTO-1449 ☐ Copies of IDS Citations
- ☐ Preliminary Amendment
- ☒ Return Receipt Postcard (MPEP 503)
(Should be specifically itemized)
- ☐ * Small Entity Statement(s) ☐ Statement filed in prior application, Status still proper and desired (PTO/SB/09-12)
- ☐ Certified Copy of Priority Document(s) (if foreign priority is claimed)
- ☐ Other:

* NOTE FOR ITEMS 1 & 14: IN ORDER TO BE ENTITLED TO PAY SMALL ENTITY FEES, A SMALL ENTITY STATEMENT IS REQUIRED (37 C.F.R. § 1.27), EXCEPT IF ONE FILED IN A PRIOR APPLICATION IS RELIED UPON (37 C.F.R. § 1.28).

17. If a CONTINUING APPLICATION, check appropriate box, and supply the requisite information below and in a preliminary amendment:
- | | | | |
|---|-------------------------------------|---|--------------------------------|
| <input type="checkbox"/> Continuation | <input type="checkbox"/> Divisional | <input type="checkbox"/> Continuation-in-part (CIP) | of prior application No: _____ |
| Prior application information: Examiner _____ | | | Group / Art Unit: _____ |

18. CORRESPONDENCE ADDRESS

<input type="checkbox"/> Customer Number or Bar Code Label	(Insert Customer No. or Attach bar code label here)	or <input checked="" type="checkbox"/> Correspondence address below
Name	Jeffrey B. Oster Davis Wright Tremaine	
Address	2600 Century Square 1501 Fourth Avenue	
City	Seattle	State WA Zip Code 98101-1688
Country	USA	Telephone 206-628-7711 Fax 206-628-7699

Name (Print/Type)	Jeffrey B. Oster	Registration No. (Attorney/Agent)	32,585
Signature	<i>Jeffrey B. Oster</i>	Date	20 January 1999

Burden Hour Statement: This form is estimated to take 0.2 hours to complete. Time will vary depending upon the needs of the individual case. Any comments on the amount of time you are required to complete this form should be sent to the Chief Information Officer, Patent and Trademark Office, Washington, DC 20231. DO NOT SEND FEES OR COMPLETED FORMS TO THIS ADDRESS. SEND TO: Assistant Commissioner for Patents, Box Patent Application, Washington, DC 20231.

HER-2 BINDING ANTAGONISTS

Technical Field of the Invention

5 The present invention provides a HER-2 binding antagonist. Specifically, intron retention has generated a novel HER-2 antagonist polypeptide that binds to the HER-2 receptor.

This work was supported by a grant from the Department of Defense (DOD) Breast Cancer Research Program. The United States Government has certain rights in this invention.

Background of the Invention

10 The HER-2/neu (erbB-2) oncogene encodes a receptor-like tyrosine kinase (RTK) that has been extensively investigated because of its role in several human carcinomas (Hynes and Stern, *Biochim. et Biophys. Acta* 1198:165-184, 1994; and Dougall et al., *Oncogene* 9:2109-2123, 1994) and in mammalian development (Lee et al., *Nature* 378:394-398, 1995). The sequence of the HER-2 protein was determined from a cDNA that was cloned by homology to the epidermal growth factor receptor (EGFR) mRNA from placenta (Coussens et al., *Science* 230:1132-1139, 15 1985) and from a gastric carcinoma cell line (Yamamoto et al., *Nature* 319:230-234, 1986). The HER-2 mRNA was shown to be about 4.5 kb (Coussens et al., *Science* 230:1132-1139, 1985; and Yamamoto et al., *Nature* 319:230-234, 1986) and encodes a transmembrane glycoprotein of 185 kDa in normal and malignant human tissues (p185HER-2) (Hynes and Stern, *Biochim. et Biophys. Acta* 1198:165-184, 1994; and Dougall et al., *Oncogene* 9:2109-2123, 1994). The function of the HER-2 gene has been examined mainly by expressing the cDNA corresponding to the 4.5 kb transcript in transfected cells and from the structure and biochemical properties of the 185 kDa protein product. P185HER-2 consists of a large extracellular domain, a transmembrane segment, and an intracellular domain with tyrosine kinase activity (Hynes and Stern, *Biochim. et Biophys. Acta* 1198:165-184, 1994; and Dougall et al., *Oncogene* 9:2109-2123, 1994). Overexpression of p185HER-2 causes phenotypic transformation of cultured cells (DiFiore et al., *Science* 237:178-182, 1987; and Hudziak et al., *Proc. Natl. Acad. Sci. USA* 84:7159-7163, 1987) and has been associated with aggressive clinical progression of breast and ovarian cancer (Slamon et al., *Science* 235:177-182, 1987; and Slamon et al., *Science* 244:707-712, 1989). p185HER-2 is highly homologous to the EGFR. However, a ligand that directly binds with high affinity to p185HER-2 has not yet been identified. Moreover, the signaling activity of HER-2 may be mediated through heterodimerization with other ligand-binding members of the EGFR family (Carraway and Cantley, *Cell* 78:5-8, 1994; Earp et al., *Breast Cancer Res. Treat.* 35:115-132, 1995; and Qian et al., *Oncogene* 10:211-219, 1995).

35 Divergent proteins, containing regions of the extracellular domains of HER family RTKs,

are generated through proteolytic processing of full length receptors (Lin and Clinton, *Oncogene* 6:639-643, 1991; Zabrecky et al., *J. Biol. Chem.* 266:1716-1720, 1991; Pupa et al., *Oncogene* 8:2917-2923, 1993; Vecchi et al., *J. Biol. Chem.* 271:18989-18995, 1996; and Vecchi and Carpenter, *J. Cell Biol.* 139:995-1003, 1997) and through alternative RNA processing (Petch et al., *Mol. Cell. Biol.* 10:2973-2982, 1990; Scott et al., *Mol. Cell. Biol.* 13:2247-2257, 1993; and Lee and Maihle, *Oncogene* 16:3243-3252, 1998). The extracellular domain of p185HER-2 is proteolytically shed from breast carcinoma cells in culture (Petch et al., *Mol. Cell. Biol.* 10:2973-2982, 1990; Scott et al., *Mol. Cell. Biol.* 13:2247-2257, 1993; and Lee and Maihle, *Oncogene* 16:3243-3252, 1998), and is found in the serum of some cancer patients (Leitzel et al., *J. Clin. Oncol.* 10:1436-1443, 1992) where it may be a serum marker of metastatic breast cancer (Leitzel et al., *J. Clin. Oncol.* 10:1436-1443, 1992) and may allow escape of HER-2-rich tumors from immunological control (Baselga et al., *J. Clin. Oncol.* 14:737-744, 1996; and Brodowicz et al., *Int. J. Cancer* 73:875-879, 1997).

A truncated extracellular domain of HER-2 is also the product of a 2.3 kb alternative transcript generated by use of a polyadenylation signal within an intron (Scott et al., *Mol. Cell. Biol.* 13:2247-2257, 1993). The alternative transcript was first identified in the gastric carcinoma cell line, MKN7 (Yamamoto et al., *Nature* 319:230-234, 1986; and Scott et al., *Mol. Cell. Biol.* 13:2247-2257, 1993) and the truncated receptor was located within the perinuclear cytoplasm rather than secreted from these tumor cells (Scott et al., *Mol. Cell. Biol.* 13:2247-2257, 1993). However, no particular therapeutic, diagnostic or research utility has been ascribed to this truncated extracellular domain polypeptide. A truncated extracellular domain of the EGFR, generated by alternative splicing (Petch et al., *Mol. Cell. Biol.* 10:2973-2982, 1990) is secreted, exhibits ligand-binding, and dimerization properties (Basu et al., *Mol. Cell. Biol.* 9:671-677, 1989), and may have a dominant negative effect on receptor function (Basu et al., *Mol. Cell. Biol.* 9:671-677, 1989; and Flickinger et al., *Mol. Cell. Biol.* 12:883-893, 1992).

Therefore, there is a need in the art to find molecules that bind to cellular HER-2 and particularly molecules that bind to different sites than humanized antibodies to HER-2 (e.g., Herceptin®). Such molecules would be useful therapeutic agents for various cancers that overexpress HER-2.

Summary of the Invention

The present invention provides an isolated polypeptide having from about 50 to 79 amino acids taken from the sequence of SEQ ID NO. 1, wherein the polypeptide binds to the extracellular domain ECD of HER-2 at an affinity of at least 10^8 . Preferably, the isolated polypeptide is from about 69 to 79 amino acids in length. Preferably, the isolated polypeptide binds to a site on the

ECD of HER-2 that is different from the site of binding of Herceptin® (a marketed humanized monoclonal antibody that is used for the treatment of cancer and that binds to the ECD or HER-2).

The present invention further provides an isolated DNA sequence that codes on expression for a polypeptide having from about 50 to 79 amino acids taken from the sequence of SEQ ID NO. 1, wherein the polypeptide binds to the extracellular domain ECD of HER-2 at an affinity of at least 10^8 . Preferably, the isolated polypeptide is from about 69 to 79 amino acids in length. Preferably, the isolated polypeptide binds to a site on the ECD of HER-2 that is different from the site of binding of Herceptin (a marketed humanized monoclonal antibody that is used for the treatment of cancer and that binds to the ECD or HER-2). The present invention further provides a transfected cell comprising an expression vector having a DNA sequence that codes on expression for a polypeptide having from about 50 to 79 amino acids taken from the sequence of SEQ ID NO. 1, wherein the polypeptide binds to the extracellular domain ECD of HER-2 at an affinity of at least 10^8 .

The present invention further provides an isolated and glycosylated polypeptide having from about 80 to 419 amino acids taken from the sequence of SEQ ID NO. 2, wherein the C terminal 79 amino acids are present, and wherein at least three N-linked glycosylation sites are present. Preferably, the isolated polypeptide is from about 350 to 419 amino acids in length and four N-linked glycosylation sites are present. Preferably, the isolated polypeptide binds to a site on the ECD of HER-2 that is different from the site of binding of Herceptin (a marketed humanized monoclonal antibody that is used for the treatment of cancer and that binds to the ECD or HER-2).

The present invention further provides an isolated DNA sequence that codes on expression for a polypeptide having from about 80 to 419 amino acids taken from the sequence of SEQ ID NO. 3, wherein the C terminal 79 amino acids are present, and wherein at least three N-linked glycosylation sites are present. Preferably, the isolated polypeptide is from about 350 to 419 amino acids in length and four N-linked glycosylation are present. The present invention further provides a transfected cell comprising an expression vector having a DNA sequence that codes on expression for a polypeptide having from about 80 to 419 amino acids taken from the sequence of SEQ ID NO. 3, wherein the C terminal 79 amino acids are present, and wherein at least three N-linked glycosylation sites are present.

The present invention provides a method for treating a solid tumor characterized by overexpression of HER-2, comprising administering an agent that binds to the extracellular domain (ECD) of HER-2, wherein the agent is selected from the group consisting of (a) an isolated polypeptide having from about 50 to 79 amino acids taken from the sequence of SEQ ID NO. 1, wherein the polypeptide binds to the extracellular domain ECD of HER-2 at an affinity of at least

10⁸, (b) an isolated and glycosylated polypeptide having from about 80 to 419 amino acids taken from the sequence of SEQ ID NO. 2, wherein the C terminal 79 amino acids are present, and wherein at least three N-linked glycosylation sites are present, (c) a monoclonal antibody that binds to the ECD of HER-2, and (d) combinations thereof, with the proviso that the agent cannot be the monoclonal antibody alone. Preferably, the solid tumor that overexpresses HER-2 is selected from the group consisting of breast cancer, small cell lung carcinoma, ovarian cancer and colon cancer. Preferably, the agent is the isolated polypeptide having from about 50 to 79 amino acids taken from the sequence of SEQ ID NO. 1. Most preferably, the agent is a combination of the isolated polypeptide having from about 50 to 79 amino acids taken from the sequence of SEQ ID NO. 1 and the monoclonal antibody that binds to the ECD of HER-2.

The present invention further provides a pharmaceutical composition for treating tumors that overexpress HER-2, comprising an agent selected from the group consisting of (a) an isolated polypeptide having from about 50 to 79 amino acids taken from the sequence of SEQ ID NO. 1, wherein the polypeptide binds to the extracellular domain ECD of HER-2 at an affinity of at least 10⁸, (b) an isolated and glycosylated polypeptide having from about 80 to 419 amino acids taken from the sequence of SEQ ID NO. 2, wherein the C terminal 79 amino acids are present, and wherein at least three N-linked glycosylation sites are present, (c) a monoclonal antibody that binds to the ECD of HER-2, and (d) combinations thereof, with the proviso that the agent cannot be the monoclonal antibody alone, and pharmaceutically acceptable carrier. Preferably, the agent is the isolated polypeptide having from about 50 to 79 amino acids taken from the sequence of SEQ ID NO. 1. Most preferably, the agent is a combination of the isolated polypeptide having from about 50 to 79 amino acids taken from the sequence of SEQ ID NO. 1 and the monoclonal antibody that binds to the ECD of HER-2.

The present invention further provides a method for targeting a therapeutic agent to solid tumor tissue, wherein the solid tumor tissue is characterized by overexpression of HER-2, comprising attaching the therapeutic agent to an isolated polypeptide having from about 50 to 79 amino acids taken from the sequence of SEQ ID NO. 1, wherein the polypeptide binds to the extracellular domain ECD of HER-2 at an affinity of at least 10⁸. Preferably, the isolated polypeptide is from about 69 to 79 amino acids in length. Preferably, the isolated polypeptide binds to a site on the ECD of HER-2 that is different from the site of binding of Herceptin® (a marketed humanized monoclonal antibody that is used for the treatment of cancer and that binds to the ECD of HER-2).

The present invention further provides a method for determining the prognosis of tumor treatment for a tumor that overexpresses HER-2, comprising: (a) obtaining a bodily fluid, wherein the bodily fluid is selected from the group consisting of blood, serum, urine, lymph, saliva, tumor

tissue, and combinations thereof; and (b) measuring the amount of p68HER-2 expressed using an anti-p68HER-2 antibody-based assay, wherein the assay is selected from the group consisting of ELISA, immunoprecipitation, immunohistochemistry, and Western analysis. Preferably, the method for determining the prognosis of tumor treatment further comprises measuring the amount of p185HER-2 ECD in the bodily fluid, and determining a ratio between the amount of p68HER-2 and p185HER-2.

Brief Description of the Drawings

Figure 1 shows a nucleotide sequence and amino acid of the insert in the extracellular domain of HER-2. The HER-2 ECD coding sequence from exon 1-9 (primers A and B) was amplified by PCR from a cDNA library from SKOV-3 cells. A product of ~1420 bp was found to be HER-2-specific by Southern blot analysis. This product was subcloned and the nucleotide sequence was determined. In panel A, the nucleotide sequence is shown for the 274 bp insert (outside the box) and for the immediately adjacent 5' and 3' sequences enclosed in the box. The insert is located between nucleotide residues 1171 and 1172 and following amino acid residue 340 in p185HER-2 using the numbering of Coussens et al. (*Science* 230:1132-1139, 1985). The consensus 5' and 3' splice sites at the arrows are shown in larger print. The inserted sequence is in-frame with 5' HER-2 exon sequence and is deduced to encode a 79 amino acid extension following Arg 340 (R³⁴⁰). The novel 79 novel amino acid sequence encoded by the insert is proline-rich (19%) and has a consensus asparagine linked glycosylation site, which is underlined. A stop codon was found at nucleotides 236-238 within the inserted sequence. In panel B, the predicted product of the alternative transcript is a truncated secreted protein which contains subdomains I and II identical to p185 and is missing the transmembrane domain and cytoplasmic domain. If fully glycosylated, the expected size is 65-70 kDa. This polypeptide product is referred to as p68HER-2. Thus, the product will be a truncated secreted protein which is missing the transmembrane domain and cytoplasmic domain found in p185HER-2.

Figure 2 shows the detection of alternative HER-2 transcripts containing the ECDIIIa sequence by Northern blot analysis. PolyA⁺ mRNA (2.5 µg) from different human fetal tissues (Clontech) or isolated from HEK-293 cells was resolved in a formalin agarose gel and transferred to a BrightStar® membrane (Ambion) in 10xSSC. The membrane was hybridized with a ³²P-labeled antisense RNA probe complimentary to the ECDIII sequence, stripped and reprobed with a ³²P labeled cDNA probe specific for the 5' HER-2 exon sequence. The membranes were washed under high stringency conditions and analyzed by phosphorimaging (Molecular Dynamics).

Figure 3 shows a sequence-specific reactivity of anti-ECDIIIa with a protein of ~ 68 kDa in a human embryonic kidney cell line (HEK293). Cell extract protein (20 µg) and 20 µl of media

conditioned by HEK-293 cells were Western blotted and probed with anti-ECDIIIa diluted 1:10,000 (lanes 1 and 2) or with anti-ECDIIIa diluted 1:10,000 containing 50 µg/ml purified His-tagged ECDIIIa peptide (lanes 3, 4).

Figure 4 shows the expression of p185HER-2, relative to p68ECDIIIa expression, is markedly elevated in carcinoma cell lines in which the HER-2 gene is amplified. Cell extracts (15 µg of protein) from human embryonic kidney cell line (HEK293), nontumorigenic ovarian surface epithelial cell line (IOSEVAN), ovarian carcinoma cell line with HER-2 gene amplification (SKOV-3), nontumorigenic breast epithelial cell line (HBL100), and breast carcinoma cell lines with HER-2 gene amplification (BT474 and SKBR-3), were resolved by SDS-PAGE in 7.5% acrylamide gels and analyzed as a Western blot. The Western blot was probed with both antibodies specific for p68HER-2 (anti-ECDIIIa) and for p185HER-2 (anti-neu(C)).

Figure 5 shows that p68ECDIIIa binds to p185HER-2. In panel A: Two mg of SKBR-3 cells extracted in nondenaturing buffer were immunoprecipitated with 5 µl anti-neu(N) specific for the N-terminal sequence of p68HER-2 and p185HER-2, or with 5 µl anti-neu(C) specific for the C-terminus of p185HER-2 and then probed as a Western blot with both anti-ECDIIIa specific for p68HER-2 and with anti-neu(C) specific for p185HER-2. In panel B: 100 µg of 17-3-1 cell extract were incubated in duplicate with 50 µl packed volume of NiNTA agarose (Qiagen) coupled to 20 µg of His-tagged ECDIIIa or to 20 µg His-tagged CREB fragment in 200 µl of wash buffer (20mM Tris pH 8.0, 300mM NaCl) at room temperature for 1 hr with shaking. The resin was then washed 4 times with 500 µl of wash buffer and proteins were eluted by incubation with 50 µl SDS-sample buffer at 100° C for 2 min. Eluted proteins were analyzed by Western blot analysis using antibodies against the C-terminus of p185HER-2, anti-neu(C). In panel C: Monolayers of ~10⁵ 3T3 cells or HER-2 transfected 17-3-1 cells in 12 well plates were washed twice with PBS and then incubated with 0.5 ml of serum-free media with 1% BSA and 39, 75, 150, and 300 nM of purified recombinant His-tagged ECDIIIa for 2 hrs at 4 °C. Cells were washed 1 time in PBS containing 1% BSA and twice in PBS and then were extracted in denaturing buffer. Equal aliquots (20 µg protein) were analyzed by western blotting with antibodies specific for ECDIIIa (anti-ECDIIIa) or, in the upper panel, with antibodies specific for p185HER-2 (anti-neu(C)).

Figure 6 shows that neither p68-rich conditioned media nor the ECDIIIa peptide stimulate tyrosine phosphorylation of p185HER-2. Monolayer cultures of ~10⁵ HER-2 transfected 17-3-1 cells were washed twice with PBS, incubated in serum-free media at 37 °C for 24 hrs, and then treated for 10 minutes with 75 or 150 µM His-tagged ECDIIIa or with 50X CM from HEK-293 cells that secrete high levels of p68 or 50X CM from SKOV-3 cells that have no detectible p68HER-2. The treated cells were extracted with denaturing buffer containing the phosphotyrosine phosphatase inhibitor vanadate (2 mM) and 20 µg/ml of cell extract protein from

each sample were analyzed by Western blot analysis with monoclonal antibodies against phosphotyrosine (Sigma). The blot was stripped by incubation at 55° C for 30 min in 62.5 mM Tris pH 6.7, 2% SDS, and 100 mM 2 -mercaptoethanol and then reprobed with anti-neu(C) specific for p185HER-2.

Figure 7 shows that p68HER-2 inhibited anchorage independent growth of tumorigenic cells. SKOV-3 ovarian carcinoma cells and HER-2 transfected 17-3-1 cells were suspended in media with 10% fetal bovine serum containing 0.3% agar (control conditions) to which was added 50X concentrated media conditioned by SKOV-3 cells (which contains no detectable p68HER-2 (-p68 CM)), or 50X concentrated media conditioned by HEK-293 cells (which contains 20 nM p68HER-2 (+p68CM)). Five times 10³ cells were plated in triplicate for each experimental condition onto a 0.5 ml layer of media containing 0.5% agarose in 12 well plates. The results shown are plotted as the mean and standard deviation of the number of colonies with more than 50 cells in triplicate wells counted at 21 days of incubation. Similar results were observed in three separate experiments.

Detailed Description of the Invention

The present invention is based upon the initial discovery of an alternative HER-2 mRNA of 4.8 kb with a 274 bp insert identified as intron 8. The retained intron is in-frame and encodes 79 amino acids [SEQ ID NO. 1] followed by a stop codon at nucleotide 236. The alternative mRNA predicts a truncated HER-2 protein that lacks the transmembrane and intracellular domains and contains 419 amino acids [SEQ ID NO. 2]; 340 residues that are identical to the N-terminus of p185HER-2 and 79 unique residues at the C-terminus [SEQ ID NO. 1]. Using specific antibodies against either the novel 79 amino acid residue C-terminal sequence [SEQ ID NO. 1] or the N-terminus of p185HER-2, a 68 kDa protein product was identified [SEQ ID NO.2]. This 68 kDa protein product is an alternative HER-2 transcript in cell extracts and in extracellular media from several cell lines. Expression of the alternative transcript was highest in a nontransfected human embryonic kidney cell line.

The results presented here show expression of alternative HER-2 mRNA, which contains an additional 274 nucleotides, probably intron 8. Consistent with this finding, an alternative transcript of ~ 4.8 kb was detected in human fetal kidney tissue and in the human embryonic kidney cell line, HEK 293. Moreover, a transcript of 2.6 kb, which is the size expected if the sequence is retained in the 2.3 kb truncated HER-2 mRNA (Yamamoto et al., *Nature* 319:230-234, 1986; and Scott et al., *Mol. Cell. Biol.* 13:2247-2257, 1993), was detected in human fetal liver tissue by Northern blot analysis using a probe specific for the inserted sequence or for the HER-2 ECD coding sequence (Figure 2). The inserted sequence introduces a termination codon and

predicts a novel 79 amino acid extension designated ECDIIIa at residue 340 of the p185HER-2 protein. The predicted protein therefore lacks the transmembrane and intracellular domains, but contains subdomains I and II of the extracellular domain of p185HER-2. As predicted, a secreted protein which contains N-terminal sequence of p185HER-2 and the C-terminal extension provided by the inclusion of the novel sequence was detected (Figures 3 and 5). The ECDIIIa protein was found to be 68 kDa which is the approximate size expected of the protein encoded by the alternative transcript if the five N-linked glycosylation sites found in subdomains I and II of p185HER-2 are glycosylated (Stern et al., *Mol. Cell. Biol.* 6:1729-1740, 1986).

The data presented herein demonstrate that p68HER-2 specifically binds to p185HER-2. The association with p185HER-2 may be conferred by the novel proline rich ECDIIIa domain rather than the N-terminal subdomains I and II of p68HER-2. While the HER-2 ECD, generated by *in vitro* deletion mutagenesis, also contains subdomains I and II, it does not associate with the extracellular domain of p185HER-2 unless engineered to enhance their proximity (Tzahar et al., *EMBO J.* 16:4938-4950, 1997; O'Rourke et al., *Proc. Natl. Acad. Sci. USA* 94:3250-3255, 1997; and Fitzpatrick et al., *FEBS Letters* 431:102-106, 1998). However, the unique ECDIIIa peptide binds with high affinity (nM concentrations) to p185HER-2 and to transfected 17-3-1 cells that overexpress p185HER-2 (Figure 5). Preferential binding of the ECDIIIa domain peptide to 17-3-1 cells indicates that secreted p68HER-2 interacts with the extracellular region of p185HER-2 at the cell surface. Therefore, p68HER-2 and fragments thereof appear to be a naturally occurring HER-2 binding protein, encoded by the HER-2 gene. In contrast to EGFR family ligands (Groenen et al., *Growth Factors* 11:235-257, 1994), p68HER-2 lacks an EGF homology domain and contains the first 340 amino acids of the receptor itself, p185HER.

Previously described putative HER-2 ligands were found to associate indirectly with p185HER-2 only in a heterodimer with an EGFR family member (Heldin and Ostman, *Cytokine Growth Factor Rev.* 7:33-40, 1996). Although it is possible that ECDIIIa binds indirectly to p185HER-2 through a coreceptor, this seems unlikely since detergent solubilized p185HER-2 was specifically and efficiently "pulled down" by immobilized ECDIIIa peptide (Figure 5B).

For all naturally occurring or engineered ligands for mammalian EGFR family members, binding is tightly coupled to stimulation of receptor dimerization and tyrosine phosphorylation (Hynes and Stern, *Biochim. et Biophys. Acta* 1198:165-184, 1994; Dougall et al., *Oncogene* 9:2109-2123, 1994; and Groenen et al., *Growth Factors* 11:235-257, 1994). Although they bind, neither p68HER-2 nor the ECDIIIa peptide was found to activate p185HER-2. Activation was assessed in two different cell lines that differ in the extent of p185HER-2 tyrosine phosphorylation, transfected 17-3-1 cells as well as SKOV-3 ovarian carcinoma cells.

Furthermore *in vitro* self-phosphorylation activity, which is enhanced in dimeric forms of

p185HER-2 (Dougall et al., *Oncogene* 9:2109-2123, 1994; and Lin et al., *J. Cell. Biochem.* 49, 290-295, 1992), was not stimulated by p68HER-2 or ECDIIIa. Similarly, the Argos protein, which is an extracellular inhibitor of the Drosophila EGF receptor and the only known antagonist of class I RTKs, did not simulate tyrosine phosphorylation of the receptor (Schweitzer et al.,
5 *Nature* 376:699-702, 1995). Likewise, Angiopoietin-2, a natural antagonist for the Tie 2 RTK, bound the endothelial receptor but failed to activate it (Maisonpierre et al., *Science* 277:55-60, 1997).

Without being bound by theory, since p68HER-2 occupies but does not activate, it could block dimerization of p185HER-2. By analogy, HER-2 ECD, when engineered to enhance its
10 binding to RTKs, prevented the formation of productive dimers required for transphosphorylation and receptor activation thereby having a dominant negative effect (O'Rourke et al., *Proc. Natl. Acad. Sci. USA* 94:3250-3255, 1997). In contrast to the HER-2 ECD, soluble p68HER-2 exhibited strong binding to p185HER-2, yet also contains subdomain I and II of the ECD. Since subdomain I may be the low affinity, promiscuous ligand binding site required for recruitment of p185HER-2
15 into heteromeric complexes (Tzahar et al., *EMBO J.* 16:4938-4950, 1997), p68HER-2 could block this site and thereby obstruct recruitment of p185HER-2 into dimers. Alternatively, p68HER-2 could compete with an uncharacterized ligand for binding to p185HER-2. The tissue-specific expression of p68HER-2 in human fetal liver and kidney may function to modulate the extent to which p185HER-2 is occupied during development of these organs. Moreover, the overexpression
20 of p185HER-2, relative to p68HER-2 in tumor cells with HER-2 gene amplification (Figure 3), could occur though a selective pressure based on overcoming the effects of a binding protein such as p68HER-2. Therefore, p68HER-2 is the first example of a naturally occurring p185HER-2 binding protein that may prevent activation of p185HER-2.

Pharmaceutical Composition

25 The present invention further provides a pharmaceutical composition for treating solid tumors that overexpress HER-2, comprising an agent selected from the group consisting of (a) an isolated polypeptide having from about 50 to 79 amino acids taken from the sequence of SEQ ID NO. 1, wherein the polypeptide binds to the extracellular domain ECD of HER-2 at an affinity of at least 10^8 , (b) an isolated and glycosylated polypeptide having from about 300 to 419 amino
30 acids taken from the sequence of SEQ ID NO. 2, wherein the C terminal 79 amino acids are present, and wherein at least three N-linked glycosylation sites are present, (c) a monoclonal antibody that binds to the ECD of HER-2, and (d) combinations thereof, with the proviso that the agent cannot be the monoclonal antibody alone, and pharmaceutically acceptable carrier. Preferably, the agent is the isolated polypeptide having from about 50 to 79 amino acids taken
35 from the sequence of SEQ ID NO. 1. Most preferably, the agent is a combination of the isolated

polypeptide having from about 50 to 79 amino acids taken from the sequence of SEQ ID NO. 1 and the monoclonal antibody that binds to the ECD of HER-2.

The inventive pharmaceutical composition, comprising either or both of the inventive polypeptides and/or monoclonal antibody, can be administered to a patient either by itself (complex or combination) or in pharmaceutical compositions where it is mixed with suitable carriers and excipients. Inventive polypeptide can be administered parenterally, such as by intravenous injection or infusion, intraperitoneal injection, subcutaneous injection, or intramuscular injection. Inventive polypeptide can be administered orally or rectally through appropriate formulation with carriers and excipients to form tablets, pills, capsules, liquids, gels, syrups, slurries, suspensions and the like. Inventive polypeptide can be administered topically, such as by skin patch, to achieve consistent systemic levels of active agent. Inventive polypeptide is formulated into topical creams, skin or mucosal patch, liquids or gels suitable to topical application to skin or mucosal membrane surfaces. Inventive polypeptide can be administered by inhaler to the respiratory tract for local or systemic treatment of cancers characterized by overexpressing HER-2.

The dosage of inventive polypeptide suitable for use with the present invention can be determined by those skilled in the art from this disclosure. Inventive polypeptide will contain an effective dosage (depending upon the route of administration and pharmacokinetics of the active agent) of inventive polypeptide and suitable pharmaceutical carriers and excipients, which are suitable for the particular route of administration of the formulation (*i.e.*, oral, parenteral, topical or by inhalation). The active inventive polypeptide is mixed into the pharmaceutical formulation by means of mixing, dissolving, granulating, dragee-making, emulsifying, encapsulating, entrapping or lyophilizing processes. The pharmaceutical formulations for parenteral administration include aqueous solutions of the inventive polypeptide in water-soluble form. Additionally, suspensions of the inventive polypeptide may be prepared as oily injection suspensions. Suitable lipophilic solvents or vehicles include fatty oils such as sesame oil, or synthetic fatty acid esters, such as ethyl oleate or triglycerides, or liposomes. Aqueous injection suspensions may contain substances which increase the viscosity of the suspension, such as sodium carboxymethyl cellulose, sorbitol, or dextran. The suspension may optionally contain stabilizers or agents to increase the solubility of the complex or combination to allow for more concentrated solutions.

Pharmaceutical formulations for oral administration can be obtained by combining the active compound with solid excipients, such as sugars (*e.g.*, lactose, sucrose, mannitol or sorbitol), cellulose preparations (*e.g.*, starch, methyl cellulose, hydroxypropylmethyl cellulose, and sodium carboxymethyl cellulose), gelatin, gums, or polyvinylpyrrolidone. In addition, a desintegrating

agent may be added, and a stabilizer may be added.

Processes for Synthesizing p68 and 79 aa C Terminal Region

Polypeptide synthesis is done by a group of standard procedures for polypeptide synthesis by sequential amino acids building through peptide synthesis equipment, following manufacturer's instructions for synthesizing peptides. Preferably, shorter polypeptides, of less than 100 amino acids, are best suited for the method of synthesis through sequential amino acid building of polypeptides. In addition, heterologous polypeptides can be expressed by transformed cells using standard recombinant DNA techniques to transform either prokaryotic or eukaryotic cells, provide appropriate growth media for their expression, and then purify the inventive polypeptide either from the media or from intracellular contents depending upon the type of cell used and its expression characteristics.

Methods for Treating Cancer with p68, 79 aa C Terminal Region, and Combinations

The present invention provides a method for treating a solid tumor characterized by overexpression of HER-2, comprising administering an agent that binds to the extracellular domain (ECD) of HER-2, wherein the agent is selected from the group consisting of (a) an isolated polypeptide having from about 50 to 79 amino acids taken from the sequence of SEQ ID NO. 1, wherein the polypeptide binds to the extracellular domain ECD of HER-2 at an affinity of at least 10^8 , (b) an isolated and glycosylated polypeptide having from about 300 to 419 amino acids taken from the sequence of SEQ ID NO. 2, wherein the C terminal 79 amino acids are present, and wherein at least three N-linked glycosylation sites are present, (c) a monoclonal antibody that binds to the ECD of HER-2, and (d) combinations thereof, with the proviso that the agent cannot be the monoclonal antibody alone. Preferably, the solid tumor that overexpresses HER-2 is selected from the group consisting of breast cancer, small cell lung carcinoma, ovarian cancer, prostate cancer, gastric carcinoma, cervical cancer, esophageal carcinoma, and colon cancer. Preferably, the agent is the isolated polypeptide having from about 50 to 79 amino acids taken from the sequence of SEQ ID NO. 1. Most preferably, the agent is a combination of the isolated polypeptide having from about 50 to 79 amino acids taken from the sequence of SEQ ID NO. 1 and the monoclonal antibody that binds to the ECD of HER-2.

The p68HER-2 polypeptide described herein was found to bind to HER-2 and prevent signal transduction through the kinase domain. Without being bound by theory, the unique ECDIIa domain mediates specific binding to p185HER-2 and the resulting interaction with p68ECDIIa prevents p185HER-2 dimerization and subsequent signal transduction. Therefore, p68HER-2 functions as a HER-2 antagonist to prevent signal transduction by preventing dimerization as a necessary prerequisite for signal transduction. Thus, the mechanism of p68HER-2 as a HER-2 antagonist is different from the mechanism of binding agents, such as the 79 amino

acid polypeptide described herein or a monoclonal antibody that binds to the ECD of HER-2. The inventive method provides that p68HER-2 inhibits tumor cell growth in tumors that overexpress HER-2 by providing a selective pressure for such tumor cells. Similarly, the HER-2 antagonists that are binding agents also inhibit tumor cell growth in tumors that overexpress HER-2 by providing selective pressure to such cells to prevent ligand binding to the ECD of HER-2 and prevent signal transduction even before potential dimerization.

Use of 79 aa C Terminal Region as a Targeting Molecule

The present invention further provides a method for targeting a therapeutic agent to solid tumor tissue, wherein the solid tumor tissue is characterized by overexpression of HER-2, comprising attaching the therapeutic agent to an isolated polypeptide having from about 50 to 79 amino acids taken from the sequence of SEQ ID NO. 1, wherein the polypeptide binds to the extracellular domain ECD of HER-2 at an affinity of at least 10^8 . Preferably, the isolated polypeptide is from about 69 to 79 amino acids in length. Preferably, the isolated polypeptide binds to a site on the ECD of HER-2 that is different from the site of binding of Herceptin® (a marketed humanized monoclonal antibody that is used for the treatment of cancer and that binds to the ECD of HER-2). It was discovered that the 79 amino acid polypeptide [SEQ ID NO. 1] exhibited surprising high affinity binding properties to the ECD of HER-2. Moreover, the site of such binding is different and unaffected by the site of binding of a marketed humanized monoclonal antibody (Herceptin®). Therefore, the high binding affinity enables the 79 amino acid polypeptide to function as a targeting molecule to tumor cells expressing HER-2.

Anti-p68 Antibody as a Diagnostic/Prognostic Agent

The p68HER-2 glycosylated polypeptide was expressed and used as an antigen for antibody production. Specifically, antibody specific for p68HER-2 was prepared by injecting rabbits with purified polyhistidine-tagged ECDIIIa peptide, which is the same as the intron encoded novel C-terminus or p68HER-2, the domain that binds with high affinity to p185HER-2. The isolated polyclonal antibody detected pM quantities of ECDIIIa peptide or of p68HER-2 with high specificity (see Figures 3 and 5). Thus, an antibody specific for p68HER-2 is useful as a diagnostic agent for detecting p68HER-2 in bodily fluids and tumor tissues using diagnostic techniques, such as ELISA, immunoprecipitations, immunohistochemistry or Western analysis.

Accordingly, the present invention further provides a method for determining the prognosis of tumor treatment for a tumor that overexpresses HER-2, comprising: (a) obtaining a bodily fluid, wherein the bodily fluid is selected from the group consisting of blood, serum, urine, lymph, saliva, tumor tissue, and combinations thereof; and (b) measuring the amount of p68HER-2 expressed using an anti-p68HER-2 antibody-based assay, wherein the assay is selected from the group consisting of ELISA, immunoprecipitation, immunohistochemistry, and Western

analysis. Preferably, the method for determining the prognosis of tumor treatment further comprises measuring the amount of p185HER-2 ECD in the bodily fluid, and determining a ratio between the amount of p68HER-2 and p185HER-2. The higher the ratio of p68HER-2:p185HER-2, the better the treatment prognosis.

5 P68HER-2 as a Therapeutic Agent

Without being bound by theory, but it appears that p68HER-2 or ECDIIIa peptide inhibits the growth of tumor cells that overexpress HER-2 by binding to p185HER-2 at the cells surface. This hypothesis was examined by testing anchorage independent growth of cells in the presence or absence of p68HER-2 using cells that depend on p185HER-2 overexpression for their malignant growth, yet have little or no detectable p68HER-2. Anchorage independent growth of cells in soft agar was used as a predictive model for tumor cytotoxicity. This is a common and predictive procedure to examine transforming activity and reflects the tumorigenic and oncogenic potential of cells (DiFore et al., *Science* 237:178-182, 1987; Hudziak et al., *Proc. Natl. Acad. Sci. USA* 84:7159-7163, 1987; and Baasner et al., *Oncogene* 13:901-911, 1996).

The effects of p68HER-2 on anchorage independent growth in soft agar was determined using SKOV-3 carcinoma cells and HER-2 transfected 17-3-1 cells, which are both tumorigenic and overexpress p185HER-2. The cells were suspended in media supplemented with fetal calf serum in the presence or absence of p68HER-2 and incubated for 21 days in a humidified incubator. Anchorage independent growth was quantitated by counting the number of colonies that contained more than 50 cells. Figure 7 shows that in the presence of p68HER-2, anchorage independent growth of both SKOV-3 cells and 17-3-1 cells was inhibited several fold. Accordingly, these data show that p68HER-2 is not just cytostatic, but cytotoxic and possibly apoptotic.

25 **Example 1**

This example provides the results from an experiment to investigate HER-2 mRNA diversity within the extracellular domain (ECD) coding sequence using polymerase chain reaction (PCR). A cDNA library from SKOV-3 cells (American Type Culture Collection (Rockville, MD) maintained in DMEM, supplemented with 10% fetal bovine serum and 0.05% gentamycin), an ovarian carcinoma cell line in which the HER-2 gene is amplified eight times (Tyson et al., *Am. J. Obstet. Gynecol.* 165:640-646, 1991) was examined using a forward primer specific for exon 1 (Tal et al., *Mol. Cell. Biol.* 7, 2597-2601, 1987) identical to nucleotides 142-161 and a reverse primer complementary to nucleotides 1265-1286 in exon 9 (Scott et al., *Mol. Cell. Biol.* 13:2247-2257, 1993). Briefly, The SKOV-3 cDNA library was provided by Origene Technologies, Inc. (Rockville, MD), and was prepared from RNA extracted from SKOV-3 cells. RNA was extracted

from SKOV-3 cells grown to 80% confluence on 15 cm plates with TriReagent (Molecular Research Center, Inc., Cincinnati, OH), according to the manufacturer's protocol, to obtain total RNA. RNA was resuspended in 10mM Tris-EDTA, pH 8.0, for reverse transcription and cDNA library construction, or in RNA hybridization buffer (80% formamide, 40mM PIPES, 4 mM NaCl, 1mM EDTA, pH 7.5) for ribonuclease protection assay (RPA). RNA concentrations were determined spectrophotometrically at OD₂₆₀. Poly A⁺ mRNA was selected from total RNA using a mRNA extraction kit (Oligotex, Qiagen).

A product of ~1420 bp, determined to be HER-2-specific by Southern blotting, was approximately 270 bp larger than the expected size of 1144 bp from the previously described cDNA sequence (Coussens et al., *Science* 230:1132-1139, 1985). Briefly, the Southern blotting procedure transferred nucleic acids from agarose gels under vacuum (Bio-Rad Model 785 Vacuum Blotter) in 0.4 M NaOH to Gene Screen Plus Hybridization Transfer Membrane (NEN Research Products, Boston, MA). Nucleic acids were fixed to membranes by UV crosslinking in a UV-Stratalinker (Stratagene, Inc., La Jolla, CA), and the membranes were blocked in hybridization buffer (50% formamide, 5X SSC, 1% SDS, 10 mg/ml herring sperm DNA) at 42 °C for 2 h. The membranes were hybridized at 42 °C for 16 h in hybridization buffer with 10⁷ cpm of a 220 bp Kpn-HincII fragment from ECDIIIa cDNA labelled with (α-³²P)dCTP (NEN Life Sciences) using a Random Prime DNA Labelling Kit (Boehringer Mannheim).

Templates were amplified in a Perkin Elmer GeneAmp PCR System 2400 (Perkin Elmer Cetus, Emeryville, CA) using the Expand High Fidelity PCR System (Boehringer Mannheim) with 1X High Fidelity PCR buffer containing 2.5 mM MgCl₂, 5 μM of each primer, and 200 μM dNTPs. All primers were obtained from GIBCO BRL (Life Technologies). Numbering of nucleotide and amino acid residues is according to the HER-2 cDNA sequence reported by Coussens et al. (Coussens et al., *Science* 230:1132-1139, 1985). The HER-2 extracellular domain was targeted for amplification from an SKOV-3 cDNA library (Origene Technologies, Inc.) using a forward primer (A) identical to nucleotides (nt) 142-161 of HER-2 cDNA (5'-TGAGCACCATGGAGCTGGC-3' [SEQ ID NO 3]), which spans the initiation codon (underlined) and a reverse primer (B) (5'-TCCGGCAGAAATGCCAGGCTCC-3' [SEQ ID NO 4]), which is complementary to HER-2 exon sequence at nt 1265-1286. Cycling parameters were: 94 °C, 30 sec; 58 °C, 45 sec; 68 °C, 3 min, for 30 cycles. The region spanning the alternative sequence (denoted ECDIIIa) from genomic DNA, was amplified using a forward primer (C) (5'-AACACAGCGGTGTGAGAAGTGC-3' [SEQ ID NO 5]) identical to HER-2 exon-specific sequence at nt 1131-1152 and the reverse primer (B) [SEQ ID NO. 4] on DNA prepared as described (Bond et al., *FEBS Letters* 367:61-66, 1995) with cycling parameters: 94 °C, 30 sec; 62 °C, 30 sec; 72 °C, 60 sec, for 25 cycles.

Reverse transcriptase-polymerase chain reaction (RT-PCR) was used to investigate the structure of mRNA containing the ECDIIIa sequence. First strand cDNA was reverse transcribed (Bond et al., *FEBS Letters* 367:61-66, 1995) using 5 µg RNA primed with 0.5 µg oligo-dT. To amplify the ECDIIIa insert and adjacent 5' HER-2 exon sequence, a forward primer (A) described above and a reverse primer (D) (5'-ATACCGGGACAGGTCAACAGC-3' [SEQ ID NO 6]) which is complementary to the 3'ECDIIIa-specific sequence were used. Cycling parameters were: 94 °C, 30 sec; 60 °C, 40 sec; 68 °C, 2 min, for 30 cycles.

Amplification of the ECDIIIa insert and adjacent 3' HER-2 exon-specific sequence was with a forward primer (E) (5'-TCTGGGTACCCACTCACTGC-3' [SEQ ID NO 7]) which is identical to the 5'ECDIIIa-specific sequence and contains a *Kpn*I restriction site and a reverse primer (F) (5'-TTCACACTGGCAGTCCAGACC-3' [SEQ ID NO 8]) which is complementary to HER-2 exon sequence at nt 3898-3919 and spans the termination codon (underlined). Cycling parameters were: 94 °C, 30 sec; 60 °C, 40 sec; 68 °C, 5 min, for 30 cycles.

The PCR product was subcloned and the nucleotide sequence was determined.

The results showed that the normal HER-2 coding sequence was present beginning with the 5' primer sequence and continued uninterrupted through nucleotide 1171. At this position, a 274 nucleotide insertion was found, followed by the expected coding sequence, including the 3' primer sequence. Analysis of the predicted protein product showed that the 274 nucleotide insertion encodes an extension of the known HER-2 protein, beginning at residue 340 (Coussens et al., *Science* 230:1132-1139, 1985), and introduces an in-frame stop codon 79 amino acids later (Figure 1). Comparison of the inserted nucleotides and their predicted amino acid sequence with sequences in Genbank showed no homologies. Examination of the 5' and 3' junctions of the divergent sequence revealed consensus splice donor and acceptor sites (Sharp, and Burge, *Cell* 91:875-879, 1997) and include a pyrimidine tract and potential branchpoint adenine residues near the 3' end of the insert sequence (Figure 1). Thus, the inserted sequence is likely to be an intron.

Inspection of the predicted amino acid sequence of the novel 79 amino acids [SEQ ID NO. 1] encoded by the inserted sequence shows a consensus N-linked glycosylation site and a high proline content of 19% (Figure 1). The inserted sequence was designated ECDIIIa since it is located at the boundary between subdomains II and III in the extracellular domain of the p185HER-2 sequence (Lax et al., *Mol. Cell. Biol.* 8:1831-1834, 1988). The insert sequence is in-frame with the adjacent 5' HER-2 exon sequence for 236 nt where there is a termination codon.

Example 2

This example provides the results from experiments characterizing ECDIIIa as contiguous with HER-2 exons in the genome. To investigate the HER-2 gene structure in the region of the

ECDIIIa sequence, a forward primer, identical to nucleotides 763-785, and a reverse primer, complementary to nucleotides 1265-1286 of the HER-2 cDNA, were used in the PCR on human genomic DNA. The amplification product was anticipated to span exon 5 (Tal et al., *Mol. Cell. Biol.* 7:2597-2601, 1987) to an exon which is immediately 3' of the ECDIIIa sequence. Intron number and sizes were estimated based on PCR product sizes, restriction digest analysis, and partial sequence analysis of amplification products.

Next, human genomic DNA was examined using HER-2 exon-specific primers that directly flank the insert to determine the sequences immediately flanking the ECDIIIa sequence. A ~430 bp product was amplified from normal human genomic DNA and from genomic DNA extracted from carcinoma cell lines SKOV-3, SKBR-3 and BT474, all of which have HER-2 gene amplification (Kraus et al., *EMBO J.* 6:605-610, 1987) and were found to express ECDIIIa in their cDNA. The identities of the PCR products as HER-2 were verified by Southern blot analysis using the procedure described in Example 1. Nucleotide sequence analysis showed that the PCR product from human genomic DNA contained the ECDIIIa insert, flanked immediately on both sides by known HER-2 coding sequence; no mutations or rearrangements were seen. These data show that the ECDIIIa sequence represents a wholly retained intron, likely intron 8 based on the size of products amplified following intron 4 and on the location of intron 8 in the homologous EGFR gene and HER-3 gene (Lee and Maible, *Oncogene* 16:3243-3252, 1998).

Example 3

This example shows that ECDIIIa is the only retained intron within the coding sequence of HER-2 mRNA. To determine whether additional introns were retained in the mRNA containing the ECDIIIa insert sequence, the reverse transcriptase-polymerase chain reaction (RT-PCR) was employed. First, a forward primer identical to 5' HER-2 cDNA sequence at 142-161 which spans the initiation codon, and a reverse primer complementary to the 3' ECDIIIa sequence were employed with SKBR-3 and SKOV-3 cDNA. A product of 1.3 kb was amplified, which is the size expected if the product contained no introns other than intron 8. Amplification of the 3'HER-2 coding sequence was then performed using a forward primer identical to 5' ECDIIIa sequence and a reverse primer complementary to 3'HER-2 cDNA sequence at nucleotides 3898-3919, which spans the p185HER-2 termination codon. A product of 2.9 kb was amplified, which is the size expected from the HER-2 cDNA if no additional introns were retained.

Further characterizations of both the 5'(1.3 kb) and 3'(2.9 kb) amplification products by restriction digest analysis and nucleotide sequencing confirmed the absence of additional retained introns. To determine the size of the products amplified when intron sequences are included, genomic DNA was used as a template for the PCR reactions, which resulted in products of

approximately 10 kb for the 5' coding sequence and 5 kb for the 3' coding sequence. These results indicate that the alternative HER-2 transcript, resulting from retention of an intron of 274 bp, was expected to be about 4.8 kb in size, assuming that the 5'untranslated (5'UTR) and 3'untranslated (3'UTR) regions are identical in size to the previously described ~4.5 kb HER-2 cDNA (Coussens et al., *Science* 230:1132-1139, 1985).

Example 4

This example illustrates the expression of a protein containing an ECDIIIa sequence. To assess whether the alternative sequence is translated into a protein product, the ECDIIIa sequence was expressed as a polyhistidine-tagged peptide in bacteria, purified the peptide by nickel-affinity chromatography, and raised antisera against the purified peptide. Briefly, the bacterial expression vector was prepared by amplifying the ECDIIIa sequence from the SKOV-3 cDNA library using primer E and a reverse primer complementary to the 3' end of the ECDIIIa insert sequence. The reverse primer contained a *Bam*H1 restriction site sequence, and was identical to that used for template construction in the RPA (described in examples 1 and 2). The PCR amplification product of ~280 bp was digested with *Kpn*I and *Bam*H1, gel purified (Qiaex II, Qiagen, Chatsworth, CA), and cloned into the pET30a vector, which encodes a six histidine tag at the amino-terminus of the expressed protein (Novagen, Madison, WI). The resulting expression vector, pET-ECDIIIa, was used for transformation of bacterial strain BL21.

To express the ECDIIIa protein product, BL21 cells transformed with the pET-ECDIIIa expression vector were grown in LB broth with 30 µg/ml Kanamycin for 4 h at 37 °C. Expression was induced with 0.1 mM IPTG for 3 h and the harvested cells were lysed by sonication, and then centrifuged at 39,000 x g for 20 min. The supernatant was absorbed onto Ni-NTA agarose (Qiagen), by shaking for 60 min at room temperature. The resin was washed with ten volumes of wash buffer (10 mM Tris pH 7.9 and 300 mM NaCl), followed by ten volumes of wash buffer with 50 mM imidazole. The his-tagged ECDIIIa protein was eluted in wash buffer with 250 mM imidazole. The his-tagged protein, which was estimated to be approximately 90% pure by Coomassie Blue staining of gels, was used to generate and characterize antibodies.

Briefly, anti-ECDIIIa antisera were produced by Cocalico Biologicals, Inc. (Reamstown, PA) by injection of two rabbits with purified polyhistidine-tagged ECDIIIa peptide (described below). Polyclonal anti-*neu* (N) was produced against a peptide identical to amino acid residues 151-165 of p185HER-2 (Lin and Clinton, *Oncogene* 6:639-643, 1991). Polyclonal anti-*neu* (C) was made against a peptide identical to the last 15 residues of the carboxy-terminus of p185HER-2 (Lin et al., *Mol. Cell. Endocrin.* 69:111-119, 1990). Antisera from two immunized rabbits were characterized and found to contain antibodies of high titer that reacted with the purified ECDIIIa

peptide.

A Western blot analysis examined whether SKBR-3 cells, which expressed the alternative sequence in its cDNA, produced a protein that reacts with anti-ECDIIIa antibody. A 68 kDa protein from the cell extract and from the extracellular media reacted with anti-ECDIIIa antibody from two different rabbits diluted at least 20,000 fold, but not with preimmune sera. Inspection of the cDNA sequence of the alternative transcript (Figure 1) predicted a secreted protein product of 65-70 kDa if all 5 consensus N-linked glycosylation sites in the N-terminal p185HER-2 sequence were glycosylated (Stern et al., *Mol. Cell. Biol.* 6:1729-1740, 1986).

If the 68 kDa ECDIIIa protein [SEQ ID NO. 2] is the translation product of the alternative HER-2 mRNA, then its N-terminal residues should be identical to the N-terminal 340 residues of p185HER-2. Therefore, cell extract from SKBR-3 cells was immunoprecipitated with anti-peptide antibody against an N-terminal sequence of HER-2, anti-neu (N) (Lin and Clinton, *Oncogene* 6:639-643, 1991) or with anti-ECDIIIa, and the immune complexes were examined by Western blot analysis with both antibodies. Briefly, three to 5 μ l of antisera were added to 2 mg of protein from cell lysates prepared in M-RIPA buffer (1% Nonidet P-40, 50 mM Tris pH 7.4, 0.1% sodium deoxycholate, 150 mM NaCl, 1 mM PMSF, 1% aprotinin), which had been centrifuged to remove nuclei. Immunoprecipitation was for 2 h with shaking at 4 °C as described (Lin et al., *Mol. Cell. Endocrin.* 69:111-119, 1990). The immune complexes were bound to Protein G Sepharose (Pharmacia) by incubation for 1 h at 4 °C with shaking, collected by centrifugation, and washed four times with M-RIPA. The proteins were released from the immune complex by incubation at 95° C for 2 min in SDS-PAGE sample buffer and resolved by SDS-PAGE in 7.5% gels (Mini-Protean II electrophoresis cell, Bio-Rad).

Western blotting was conducted following SDS-PAGE. Proteins were electroblotted onto nitrocellulose (Trans-blot, BioRad) using a semi-dry transfer unit (Bio-Rad) at 15 V for 20 min per gel (0.75 mm thick) equilibrated with 25 mM Tris pH 8.3, 192 mM glycine, 50 mM NaCl, and 20% methanol. The membranes were blocked with 5% nonfat dry milk at 25 °C for one hour. The blots were then incubated with primary antibody, washed twice for 15 min, and four times for 5 min with TBS-Tween (Tris-buffered saline containing 0.05% Tween), and then incubated for 40 min with goat anti-rabbit secondary antibody, conjugated to horseradish peroxidase (Bio-Rad), diluted 1:10, 000 in TBS-Tween. After incubation with secondary antibody, the membranes were washed as described above and reacted with chemiluminescent reagent (Pierce) and then were exposed to Kodak X-OMAT BLU film.

As expected, p68HER-2 was detected when anti-ECDIIIa was used for immunoprecipitation and for Western blot analysis. When anti-ECDIIIa was used for immunoprecipitation and anti-neu (N) was the probe in the Western blot, a 68kDa protein was

detected, indicating that p68ECDIIIa contained the N-terminal sequence of p185HER-2. Further, anti-neu (N) precipitated p68HER-2, which was detected by probing with anti-ECDIIIa antibody. These results demonstrate that p68HER-2 contains both ECDIIIa and the N-terminal sequence of HER-2.

Several other cell lines were examined for expression of p68ECDIIIa. The carcinoma cell lines which contained ECDIIIa sequence in their cDNA (BT474, SKOV-3) also had p68HER-2. Of several cell lines examined, HEK293 cells, derived from normal human embryonic kidney cells, expressed the highest levels of p68ECDIIIa in the cell extract and in the extracellular media, at about 5 to 10-fold higher amounts than SKBR-3 cells. In comparison to the carcinoma cell lines examined (SKBR-3, SKOV-3, and BT474) which overexpress p185HER-2, the HEK293 cells contained about 20 fold lower amounts of p185HER-2. Therefore, the relative proportion of p68HER-2 to p185HER-2 was at least 100 fold greater in HEK293 cells than in the three carcinoma cell lines studied. Reactivity with p68HER-2 as well as with a protein of ~120 kDa, particularly apparent in the HEK293 extracts, was blocked by preincubation of the antisera with purified ECDIIIa peptide demonstrating sequence-specific reactivity. The larger protein may be a dimer of p68HER-2. Therefore, p68HER-2 was expressed and secreted from several carcinoma cell lines and is at 5-10 fold elevated levels in HEK293.

Example 5

This example illustrates expression of an alternative HER-2 transcript containing the ECDIIIa intron sequence. Results of the RT-PCR analysis indicated that the ECDIIIa sequence was inserted into an otherwise normal-sized HER-2 mRNA. These data suggest an alternative transcript of ~4.8 kb. To examine the size and expression of the ECDIIIa alternative transcript, Northern blot analysis was conducted using an ECDIIIa-specific probe. Briefly, a template for antisense RNA probe synthesis was constructed from SKOV-3 cDNA by PCR amplification of a 389 bp sequence spanning the entire ECDIIIa insert sequence and containing adjacent 5'HER-2 exon sequence. The PCR was done using the forward primer C [SEQ ID NO. 5] that is identical to HER-2 cDNA sequence at nt 1131-1152 and a reverse primer (5'-GCACGGATCCATAGCAGACTGAG GAGG-3' [SEQ ID NO. 9]) which contains a 3' *Bam*H1 restriction endonuclease site and is complementary to the sequence spanning the 3' splice site of the ECDIIIa sequence. The PCR product was then digested with *Bam*H1, liberating a 375 bp fragment, which was cloned into pBluescript SK (Stratagene). The plasmid was sequenced by the Vollum Institute Core Sequencing Facility (Portland, OR) with m13 forward and reverse primers. An antisense RNA probe complimentary to the entire ECDIIIa sequence and to 87 nt of HER-2 exon sequence 5' to the insert was transcribed from 1 µg of linearized template using (α -³²P) CTP,

T7 RNA polymerase, and the T7/SP6 Riboprobe Synthesis System (Promega, Madison, WI). This probe was expected to protect a 370 nt fragment when hybridized with mRNA containing ECDIIIa and adjacent HER-2 exon sequence, and to protect an 87 nt fragment when hybridized with fully spliced HER-2 mRNA.

To prepare the RNA hybrids, 30 µg of RNA were hybridized with approximately 50,000 cpm of antisense RNA probe at 48 °C for 16 h. RNA hybrids were digested for 30 min at 37 °C with 40 µg/ml RNaseA (Boehringer Mannheim) and 2 µg/ml RNase T1 (Life Technologies) in a solution of 250 mM NaCl, 5 mM EDTA, and 10 mM Tris pH 7.5. Proteinase K (100 µg) (Life Technologies) in 20 µl 10% SDS was added to stop the digestion. Samples were extracted with acid phenol (pH 4.5; Life Technologies) and chloroform, precipitated with two volumes of 100% ethanol, and suspended in 5 µl of RPA sample buffer (88% formamide, 10 mM EDTA pH 8.0, 1 mg/ml xylene cyanol, and 1 mg/ml bromophenol blue). Samples were denatured at 95° C for 10 min and electrophoresed on a 5% polyacrylamide/urea gel in TBE (89 mM Tris, 89 mM borate, 2 mM EDTA pH 8.3). Gels were dried under vacuum and subjected to phosphorimager analysis for quantitation of the protected fragments (IP Lab Gel, Molecular Dynamics).

An alternative transcript of approximately 4.8 kb was detected in HEK293 cells which expressed the highest levels of p68ECDIIIa. However an alternative transcript could not be detected by Northern analysis of the SKBR-3, BT474, or SKOV-3 carcinoma cell lines. Therefore, the more sensitive ribonuclease protection assay (RPA) was employed to examine the expression levels of the alternative transcript relative to the fully spliced 4.5 kb transcript. RNA from ovarian (SKOV-3) and breast (SKBR-3 and BT474) carcinoma cell lines, which contained detectable levels of p68ECDIIIa, and a control cell line, 17-3-1, stably transfected with HER-2 cDNA, were hybridized with an antisense ³²P-labeled RNA probe which spanned the entire ECDIIIa (intron 8) sequence and 5' HER-2 exon sequence flanking intron 8. Following RNase digestion, electrophoresis, and autoradiography, a band of 370 nucleotides was detected in each cell line except for 17-3-1, which corresponds to the expected size protected by an ECDIIIa-containing HER-2 mRNA. In addition, an 87 nucleotide protected fragment was detected in all cells and is the size expected for the fully-spliced HER-2 message which is overexpressed by more than 100 fold in these carcinoma cell lines compared to normal control cell lines (Kraus et al., *EMBO J.* 6:605-610, 1987). The amounts of each protected fragment were quantitated and normalized for size to estimate the relative abundance of the alternative transcript, expressed as a percentage of the p185HER-2 mRNA. The alternative HER-2 mRNA with the ECDIIIa insert was at 4.2% the level of the fully spliced transcript in SKOV-3; 5.4% in SKBR-3, and 0.8% in BT474 cells.

Example 6

This example shows that alternative transcripts containing the ECDIIIa insert were expressed in human embryonic kidney and liver. A Northern blot was conducted to examine whether an alternative transcript, which contains the ECDIIIa sequence, was expressed in normal human tissue. PolyA⁺ mRNA from a variety of human fetal tissues prepared as a Northern blot was hybridized with a radiolabeled probe specific for the unique ECDIIIa sequence. A 4.8 kb mRNA was detected in kidney and a 2.6 kb transcript was detected in liver (Figure 2). The 4.8 kb transcript likely corresponded to the full length 4.5 kb transcript with the 274bp insert and the 2.6 kb transcript may have corresponded to a previously described 2.3 kb alternative transcript (Yamamoto et al., *Nature* 319:230-234, 1986; and Scott et al., *Mol. Cell. Biol.* 13:2247-2257, 1993) with the 274bp ECDIIIa insert. When the blot was stripped and hybridized with a probe specific for the 5' HER-2 coding sequence, a broad band representing the 4.8 and 4.5 kb mRNAs was detected in fetal kidney tissues and the truncated 2.6 kb transcript was detected in liver showing that these alternative transcripts contain sequences that encode the HER-2 ECD. Because the inserted ECDIIIa sequence contained a termination codon, the same protein product may be produced from each of these mRNAs.

Several cell lines were also investigated for the ECDIIIa-containing alternative transcript by Northern blot analysis. The 4.8 kb alternative transcript was detected in the human embryonic kidney cell line, HEK-293 (Figure 2). Although the ECDIIIa sequence was detected by RT-PCR analysis of SKBR-3, BT474, and SKOV-3 carcinoma cell lines, which all contain HER-2 gene amplification, an ECDIIIa-containing alternative transcript could not be detected by Northern analysis of these cells. Therefore, the more sensitive ribonuclease protection assay (RPA) was employed using an antisense probe which spanned the entire ECDIIIa sequence and 5' HER-2 exon sequence flanking the ECDIIIa sequence. The alternative HER-2 mRNA with the ECDIIIa insert was detected at less than 5% of the fully spliced transcript in SKOV-3, SKBR-3, and BT474 cells. These findings show that two alternative transcripts containing the ECDIIIa sequence were expressed in a tissue-specific manner in normal human tissues, that the 4.8 kb alternative transcript was expressed in the HEK-293 cell line, and that the carcinoma cells with gene amplification express reduced amounts of the alternative transcript at less than 5% of the 4.5 kb HER-2 transcript.

Example 7

This example illustrates expression of a protein containing the ECDIIIa sequence. To assess whether the alternative sequence was translated into a protein product, the ECDIIIa sequence, as a polyhistidine-tagged peptide in bacteria, was expressed and purified by nickel-

affinity chromatography, and raised antisera against the purified peptide. The HEK-293 cells, which expressed the 4.8 kb ECDIIIa alternative transcript, were examined for expression of an ECDIIIa-containing protein by Western analysis. A 68 kDa protein from the cell extract and from the extracellular media reacted with the anti-ECDIIIa antibody (Figure 3) but not with preimmune sera and reactivity was blocked by preincubation of the antisera with purified ECDIIIa peptide (Figure 3). The larger protein of ~125 kDa detected in some cell extracts may be an aggregate of p68HER-2. The cDNA sequence of the alternative transcript (Figure 1) predicts a secreted protein product of 65-70 kDa if all 5 consensus N-linked glycosylation sites in the N-terminal p185HER-2 sequence are glycosylated (Stern et al., *Mol. Cell. Biol.* 6:1729-1740, 1986). Several other cell lines were examined for expression of p68ECDIIIa. The carcinoma cell lines which contained ECDIIIa sequence in their cDNA (BT474, SKOV-3, SKBR-3) also had detectable levels of p68HER-2.

Example 8

This example illustrates the expression of p68HER-2 relative to p185HER-2 was markedly reduced in carcinoma cell lines in which the HER-2 gene is amplified. Because the p68HER-2 mRNA was expressed at very low levels relative to the p185HER-2 mRNA in carcinoma cell lines with HER-2 gene amplification, the relative proportions of p68HER-2 and p185HER-2 proteins in several cell lines were examined with and without HER-2 gene amplification. Western blots were prepared and probed with both antisera specific for p68HER-2 and for p185HER-2. Figure 4 shows that p185HER-2 was readily detected in the carcinoma cells lines that have their HER-2 gene amplified about 8 times (Kraus et al., *EMBO J.* 6:605-610, 1987). However, there was not a corresponding elevation in p68HER-2. In comparison, p68HER-2 was the only HER-2 protein detected in the HEK-293, IOSEVAN, and HBL100 nontumorigenic cells, although p185HER-2 was expressed at very low levels in these cells (Kraus et al., *EMBO J.* 6:605-610, 1987) and was detected in overexposed blots. These data show that p68HER-2 was low in proportion to p185HER-2 in carcinoma cells with HER-2 gene amplification and suggests that a mechanism may exist to maintain low levels of p68HER-2 when p185HER-2 is overexpressed.

Example 9

This example illustrates that p68HER-2 and the ECDIIIa peptide specifically bind to p185HER-2. Because p68HER-2 is secreted and contains subdomains I and II identical to p185HER-2, in addition to a novel sequence, the possibility that p68HER-2 may interact with p185HER-2 was investigated. Antipeptide antibody against the N-terminus of p185HER-2 and p68HER-2, anti-neu (N), or antibody specific for p185HER-2, anti-neu(C), were used for

immunoprecipitations of SKBR-3 carcinoma cells, which express low levels of p68HER-2 and overexpress p185HER-2. The immunoprecipitated material was prepared as a Western blot and probed with both anti-ECDIIIa specific for p68HER-2 and with anti-neu(C). Anti-neu (N) immunoprecipitated both p68HER-2 and p185HER-2 (Figure 5A). In addition, antibodies specific for the C-terminus of p185HER-2 immunoprecipitated p185HER-2 and coprecipitated p68HER-2 (Figure 5A), suggesting an interaction between the two proteins.

Since binding interactions between ECD sequences are very weak (Tzahar et al., *EMBO J.* 16:4938-4950, 1997; Fitzpatrick et al., *FEBS Letters* 431:102-106, 1998), the possibility that binding may be conferred by the novel proline rich ECDIIIa domain was examined. The unique 79 amino acid domain, purified as a His-tagged protein, was immobilized on nickel agarose and used in a pull-down assay. For controls, two purified His-tagged peptides unrelated to ECDIIIa, a 600 residue fragment of the Wilson's disease membrane protein, and a 70 residue fragment containing the DNA binding domain of the CREB protein, were likewise immobilized on nickel agarose resin. The immobilized peptides were incubated with protein extracts prepared from HER-2 transfected 3T3 cells (17-3-1). Following extensive washes, the bound proteins were eluted and prepared as a Western blot which was probed with an antibody specific for p185HER-2.

Equal amounts of His-tagged ECDIIIa peptide and control peptide were bound to the resin as confirmed by elution with 1M imidazole and Coomassie staining of the eluted material in SDS-gels. While no p185HER-2 was retained by resin without peptide or with control peptide, p185HER-2 was selectively retained by the ECDIIIa peptide (Figure 5B).

Since the ECDIIIa domain bound to p185HER-2 in a pulldown assay, the question of whether the ECDIIIa domain preferentially binds to cells that overexpress p185HER-2 was examined. This was investigated using monolayer cultures of 17-3-1 cells transfected with HER-2 compared to the parental 3T3 cells. The cells were incubated with different concentrations of the His-ECDIIIa peptide, washed, and extracted in denaturing buffer with protease inhibitors. To detect any bound peptide, the cell extracts were examined by Western blot analysis using antibodies specific for ECDIIIa. In addition, equal aliquots of the ECDIIIa peptide treated cells were reacted as a Western blot with antibodies specific for p185HER-2, demonstrating the overexpression of p185HER-2 in the transfected 17-3-1 cells. The ECDIIIa peptide preferentially bound to intact 17-3-1 cells at nM concentrations (Figure 5C) whereas little or no peptide was found to bind to equivalent amounts of parental 3T3 cells suggesting a specific interaction with the extracellular domain of p185HER-2.

Example 10

Effect of p68ECDIIIa and the ECDIIIa peptide on tyrosine phosphorylation of p185HER-2

was examined. Tyrosine phosphorylation of RTKs is the initial indication of ligand activation and signal transduction. Tyrosine phosphorylation in 17-3-1 cells treated with different amounts of the purified ECDIIIa peptide, with conditioned media (CM) from HEK293 cells that contained high levels of p68HER-2 (Figure 2A), or with control, conditioned media from SKOV-3 cells that had no detectable p68HER-2 were examined. There was no increase in the tyrosine phosphorylation signal at 10 minutes (Figure 6) or 2 hrs of treatment with His-ECDIIIa or with concentrated CM suggesting that p185HER-2 was not activated. Neither p68HER-2-containing CM nor the ECDIIIa peptide detectably altered the phosphotyrosine signal corresponding to p185HER-2 from SKOV-3 cells in which p185HER-2 tyrosine phosphorylation levels were low. Additionally, p68HER-2 and the ECDIIIa peptide had no discernable effect on *in vitro* self-phosphorylation activity of p185HER-2 immunoprecipitated from 17-3-1 cell extracts. These results support the conclusion that p68HER-2 did not activate p185HER-2 signal transduction.

SEQUENCE LISTING

(1) GENERAL INFORMATION:

(i) APPLICANTS: Clinton, Gail and Doherty, Joni Kristin

(ii) TITLE OF INVENTION: HER-2 BINDING ANTAGONISTS

(iii) NUMBER OF SEQUENCES: 9

(iv) CORRESPONDENCE ADDRESS:

(A) ADDRESSEE: DAVIS WRIGHT TREMAINE

(B) STREET: 1501 Fourth Avenue, 2600 Century Square

(C) CITY: Seattle

(D) STATE: Washington

(E) COUNTRY: U.S.A.

(F) ZIP: 98101

(v) COMPUTER READABLE FORM:

(A) MEDIUM TYPE: Floppy disk

(B) COMPUTER: PC compatible

(C) OPERATING SYSTEM: Windows95

(D) SOFTWARE: Word

(vi) CURRENT APPLICATION DATA:

(A) APPLICATION NUMBER: to be assigned

(B) FILING DATE: 19 January 1999

(C) CLASSIFICATION:

(viii) ATTORNEY/AGENT INFORMATION:

(A) NAME: Oster, Jeffrey B.

(B) REGISTRATION NUMBER: 32,585

(C) REFERENCE/DOCKET NUMBER: OHSU-1

(ix) TELECOMMUNICATION INFORMATION:

(A) TELEPHONE: 206 628 7711

(B) TELEFAX: 206 628 7699

(2) INFORMATION FOR SEQ ID NO:1:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 79

(B) TYPE: polypeptide

(C) STRANDEDNESS: single

(D) TOPOLOGY: unknown

(ii) MOLECULE TYPE: HER-2 ECD antagonist

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:1:

	Gly	Thr	His	Ser	Leu	Leu	Pro	Arg	Pro	Ala	Ala	Val	Pro	Val	Pro	Leu
					5					10					15	
5	Arg	Met	Gln	Pro	Gly	Pro	Ala	His	Pro	Val	Leu	Ser	Phe	Leu	Arg	Pro
			20					25					30			
	Ser	Trp	Asp	Leu	Val	Ser	Ala	Phe	Tyr	Ser	Leu	Pro	Leu	Ala	Pro	Leu
			35					40					45			
	Ser	Pro	Thr	Ser	Val	Pro	Ile	Ser	Pro	Val	Ser	Val	Gly	Arg	Gly	Pro
		50					55					60				
10	Asp	Pro	Asp	Ala	His	Val	Ala	Val	Asn	Leu	Ser	Arg	Tyr	Glu	Gly	79
	65					70					75					

(2) INFORMATION FOR SEQ ID NO:2:

15 (i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 419

(B) TYPE: protein

(C) STRANDEDNESS: single

(D) TOPOLOGY: unknown

20 (ii) MOLECULE TYPE: polypeptide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:2:

	Met	Glu	Leu	Ala	Ala	Leu	Cys	Arg	Trp	Gly	Leu	Leu	Leu	Ala	Leu	Leu
					5					10					15	
25	Pro	Pro	Gly	Ala	Ala	Ser	Thr	Gln	Val	Cys	Thr	Gly	Thr	Asp	Met	Lys
					20					25				30		
	Leu	Arg	Leu	Pro	Ala	Ser	Pro	Glu	Thr	His	Leu	Asp	Met	Leu	Arg	His
					35					40				45		
30	Leu	Tyr	Gln	Gly	Cys	Gln	Val	Val	Gln	Gly	Asn	Leu	Glu	Leu	Thr	Tyr
			50					55					60			
	Leu	Pro	Thr	Asn	Ala	Ser	Leu	Ser	Phe	Leu	Gln	Asp	Ile	Gln	Glu	Val
		65					70					75				
	Gln	Gly	Tyr	Val	Leu	Ile	Ala	His	Asn	Gln	Val	Arg	Gln	Val	Pro	Leu
	80					85				90					95	
35	Gln	Arg	Leu	Arg	Ile	Val	Arg	Gly	Thr	Gln	Leu	Phe	Glu	Asp	Asn	Tyr
					100					105					110	
	Ala	Leu	Ala	Val	Leu	Asp	Asn	Gly	Asp	Pro	Leu	Asn	Asn	Thr	Thr	Pro
				115					120					125		
	Val	Thr	Gly	Ala	Ser	Pro	Gly	Gly	Leu	Arg	Glu	Leu	Gln	Leu	Arg	Ser
40			130					135					140			
	Leu	Thr	Glu	Ile	Leu	Lys	Gly	Gly	Val	Leu	Ile	Gln	Arg	Asn	Pro	Gln
			145				150					155				
	Leu	Cys	Tyr	Gln	Asp	Thr	Ile	Leu	Trp	Lys	Asp	Ile	Phe	His	Lys	Asn
	160					165					170					175
45	Asn	Gln	Leu	Ala	Leu	Thr	Leu	Ile	Asp	Thr	Asn	Arg	Ser	Arg	Ala	Cys
					180					185					190	

60
50
40
30
20
10
0

	His	Pro	Cys	Ser	Pro	Met	Cys	Lys	Gly	Ser	Arg	Cys	Trp	Gly	Glu	Ser	
				195					200					205			
	Ser	Glu	Asp	Cys	Gln	Ser	Leu	Thr	Arg	Thr	Val	Cys	Ala	Gly	Gly	Cys	
			210					215					220				
5	Ala	Arg	Cys	Lys	Gly	Pro	Leu	Pro	Thr	Asp	Cys	Cys	His	Glu	Gln	Cys	
		225					230					235					
	Ala	Ala	Gly	Cys	Thr	Gly	Pro	Lys	His	Ser	Asp	Cys	Leu	Ala	Cys	Leu	
	240					245					250					255	
	His	Phe	Asn	His	Ser	Gly	Ile	Cys	Glu	Leu	His	Cys	Pro	Ala	Leu	Val	
10				260						265					270		
	Thr	Tyr	Asn	Thr	Asp	Thr	Phe	Glu	Ser	Met	Pro	Asn	Pro	Glu	Gly	Arg	
			275						280					285			
	Tyr	Thr	Phe	Gly	Ala	Ser	Cys	Val	Thr	Ala	Cys	Pro	Tyr	Asn	Tyr	Leu	
			290					295					300				
15	Ser	Thr	Asp	Val	Gly	Ser	Cys	Thr	Leu	Val	Cys	Pro	Leu	His	Asn	Gln	
		305					310					315					
	Glu	Val	Thr	Ala	Glu	Asp	Gly	Thr	Gln	Arg	Cys	Glu	Lys	Cys	Ser	Lys	
	320					325					330					335	
	Pro	Cys	Ala	Arg	Gly	Thr	His	Ser	Leu	Leu	Pro	Arg	Pro	Ala	Ala		
20				340					345					350			
	Val	Pro	Val	Pro	Leu	Arg	Met	Gln	Pro	Gly	Pro	Ala	His	Pro	Val	Leu	
				355					360					365			
	Ser	Phe	Leu	Arg	Pro	Ser	Trp	Asp	Leu	Val	Ser	Ala	Phe	Tyr	Ser	Leu	
			370					375					380				
25	Pro	Leu	Ala	Pro	Leu	Ser	Pro	Thr	Ser	Val	Pro	Ile	Ser	Pro	Val	Ser	
		385					390					395					
	Val	Gly	Arg	Gly	Pro	Asp	Pro	Asp	Ala	His	Val	Ala	Val	Asn	Leu	Ser	
	400					405					410					415	
	Arg	Tyr	Glu	Gly		419											

(2) INFORMATION FOR SEQ ID NO:3:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 19

(B) TYPE: nucleic acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: unknown

(ii) MOLECULE TYPE: oligonucleotide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:3:

TGAGCACCAT GGAGCTGGC 19

(2) INFORMATION FOR SEQ ID NO:4:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 22

(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: unknown
(ii) MOLECULE TYPE: oligonucleotide
(xi) SEQUENCE DESCRIPTION: SEQ ID NO:4:

TCCGGCAGAA ATGCCAGGCT CC 22

(2) INFORMATION FOR SEQ ID NO:5:

(i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 22
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: unknown
(ii) MOLECULE TYPE: oligonucleotide
(xi) SEQUENCE DESCRIPTION: SEQ ID NO:5:

AACACAGCGG TGTGAGAAGT GC 22

(2) INFORMATION FOR SEQ ID NO:6:

(i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 21
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: unknown
(ii) MOLECULE TYPE: oligonucleotide
(xi) SEQUENCE DESCRIPTION: SEQ ID NO:6:

ATACCGGGAC AGGTCAACAG C 21

(2) INFORMATION FOR SEQ ID NO:7:

(i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 20
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: unknown
(ii) MOLECULE TYPE: oligonucleotide
(xi) SEQUENCE DESCRIPTION: SEQ ID NO:7:

TCTGGGTACC CACTCACTGC 20

(2) INFORMATION FOR SEQ ID NO:8:

(i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 22

- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: unknown
- (ii) MOLECULE TYPE: oligonucleotide
- (xi) SEQUENCE DESCRIPTION: SEQ ID NO:8:

5

TTCACACTGG CACGTCCAGA CC 22

(2) INFORMATION FOR SEQ ID NO:9:

10

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 27
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: unknown

15

(ii) MOLECULE TYPE: oligonucleotide

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:9:

GCACGGATCC ATAGCAGACT GAGGAGG 27

We claim:

1. An isolated polypeptide having from about 50 to 79 amino acids taken from the sequence of SEQ ID NO. 1, wherein the polypeptide binds to the extracellular domain ECD of HER-2 at an affinity of at least 10^8 .

2. The isolated polypeptide of claim 1, wherein the isolated polypeptide is from about 69 to 79 amino acids in length.

3. The isolated polypeptide of claim 1, wherein the isolated polypeptide binds to a site on the ECD of HER-2 that is different from the site of binding of Herceptin (a marketed humanized monoclonal antibody that is used for the treatment of cancer and that binds to the ECD or HER-2).

4. An isolated DNA sequence that codes on expression for a polypeptide having from about 50 to 79 amino acids taken from the sequence of SEQ ID NO. 1, wherein the polypeptide binds to the extracellular domain ECD of HER-2 at an affinity of at least 10^8 .

5. The isolated DNA sequence that codes on expression for a polypeptide of claim 4 wherein the isolated polypeptide is from about 69 to 79 amino acids in length.

6. The isolated DNA sequence of claim 4, wherein the isolated polypeptide binds to a site on the ECD of HER-2 that is different from the site of binding of Herceptin®.

7. A transfected cell comprising an expression vector having a DNA sequence that codes on expression for a polypeptide having from about 50 to 79 amino acids taken from the sequence of SEQ ID NO. 1, wherein the polypeptide binds to the extracellular domain ECD of HER-2 at an affinity of at least 10^8 .

8. An isolated and glycosylated polypeptide having from about 300 to 419 amino acids taken from the sequence of SEQ ID NO. 2, wherein the C terminal 79 amino acids are present, and wherein at least three N-linked glycosylation sites are present.

9. The isolated and glycosylated polypeptide of claim 6, wherein the isolated polypeptide is from about 350 to 419 amino acids in length and four N-linked glycosylation are present.

10. The isolated and glycosylated polypeptide of claim 6, wherein the isolated polypeptide binds to a site on the ECD of HER-2 that is different from the site of binding of Herceptin®.

11. An isolated DNA sequence that codes on expression for a polypeptide having from about 800 to 419 amino acids taken from the sequence of SEQ ID NO. 2, wherein the C terminal 79 amino acids are present, and wherein at least three N-linked glycosylation sites are present.

12. The isolated DNA sequence that codes on expression for a polypeptide of claim 11, wherein the isolated polypeptide is from about 350 to 419 amino acids in length and four N-linked

glycosylation sites are present.

13. A transfected cell comprising an expression vector having a DNA sequence that codes on expression for a polypeptide having from about 80 to 419 amino acids taken from the sequence of SEQ ID NO. 2, wherein the C terminal 79 amino acids are present, and wherein at least three N-linked glycosylation sites are present.

14. A method for treating a solid tumor characterized by overexpression of HER-2, comprising administering an agent that binds to the extracellular domain (ECD) of HER-2, wherein the agent is selected from the group consisting of (a) an isolated polypeptide having from about 50 to 79 amino acids taken from the sequence of SEQ ID NO. 1, wherein the polypeptide binds to the extracellular domain ECD of HER-2 at an affinity of at least 10^8 , (b) an isolated and glycosylated polypeptide having from about 80 to 419 amino acids taken from the sequence of SEQ ID NO. 2, wherein the C terminal 79 amino acids are present, and wherein at least three N-linked glycosylation sites are present, (c) a monoclonal antibody that binds to the ECD of HER-2, and (d) combinations thereof, with the proviso that the agent cannot be the monoclonal antibody alone.

15. The method of claim 14, wherein the solid tumor that overexpresses HER-2 is selected from the group consisting of breast cancer, small cell lung carcinoma, ovarian cancer and colon cancer.

16. The method of claim 14, wherein the agent is the isolated polypeptide having from about 50 to 79 amino acids taken from the sequence of SEQ ID NO. 1.

17. The method of claim 16, wherein the agent is a combination of the isolated polypeptide having from about 50 to 79 amino acids taken from the sequence of SEQ ID NO. 1 and the monoclonal antibody that binds to the ECD of HER-2.

18. A pharmaceutical composition for treating solid tumors that overexpress HER-2, comprising an agent selected from the group consisting of (a) an isolated polypeptide having from about 50 to 79 amino acids taken from the sequence of SEQ ID NO. 1, wherein the polypeptide binds to the extracellular domain ECD of HER-2 at an affinity of at least 10^8 , (b) an isolated and glycosylated polypeptide having from about 80 to 419 amino acids taken from the sequence of SEQ ID NO. 2, wherein the C terminal 79 amino acids are present, and wherein at least three N-linked glycosylation sites are present, (c) a monoclonal antibody that binds to the ECD of HER-2, and (d) combinations thereof, with the proviso that the agent cannot be the monoclonal antibody alone, and pharmaceutically acceptable carrier.

19. The pharmaceutical composition for treating solid tumors that overexpress HER-2 of claim 18, wherein the agent is the isolated polypeptide having from about 50 to 79 amino acids taken from the sequence of SEQ ID NO. 1.

20. The pharmaceutical composition for treating solid tumors that overexpress HER-2 of claim 19, wherein the agent is a combination of the isolated polypeptide having from about 50 to 79 amino acids taken from the sequence of SEQ ID NO. 1 and the monoclonal antibody that binds to the ECD of HER-2.

21. A method for targeting a therapeutic agent to solid tumor tissue, wherein the solid tumor tissue is characterized by overexpression of HER-2, comprising attaching the therapeutic agent to an isolated polypeptide having from about 50 to 79 amino acids taken from the sequence of SEQ ID NO. 1, wherein the polypeptide binds to the extracellular domain ECD of HER-2 at an affinity of at least 10^8 .

22. The method for targeting a therapeutic agent to solid tumor tissue of claim 21, wherein the isolated polypeptide is from about 69 to 79 amino acids in length.

23. The method for targeting a therapeutic agent to solid tumor tissue of claim 221, wherein the isolated polypeptide binds to a site on the ECD of HER-2 that is different from the site of binding of Herceptin®.

24. A method for determining the prognosis of tumor treatment for a tumor that overexpresses HER-2, comprising: (a) obtaining a bodily fluid, wherein the bodily fluid is selected from the group consisting of blood, serum, urine, lymph, saliva, tumor tissue, and combinations thereof; and (b) measuring the amount of p68HER-2 expressed using an anti-p68HER-2 antibody-based assay, wherein the assay is selected from the group consisting of ELISA, immunoprecipitation, immunohistochemistry, and Western analysis.


25. The method for determining the prognosis of tumor treatment for a tumor that overexpresses HER-2 of claim 24, further comprising measuring the amount of p185HER-2 ECD in the bodily fluid.

26. The method for determining the prognosis of tumor treatment for a tumor that overexpresses HER-2 of claim 24, further comprising determining a ratio between the amount of p68HER-2 and p185HER-2, whereby the higher the p68HER-2 to p185HER-2 ratio, the better the prognosis of the patient.


Abstract

There is disclosed a pharmaceutical composition for treating solid tumors that overexpress HER-2, comprising an agent selected from the group consisting of (a) an isolated polypeptide having from about 50 to 79 amino acids taken from the sequence of SEQ ID NO. 1, wherein the polypeptide binds to the extracellular domain ECD of HER-2 at an affinity of at least 10^8 , (b) an isolated and glycosylated polypeptide having from about 300 to 419 amino acids taken from the sequence of SEQ ID NO. 2, wherein the C terminal 79 amino acids are present, and wherein at least three N-linked glycosylation sites are present, (c) a monoclonal antibody that binds to the ECD of HER-2, and (d) combinations thereof, with the proviso that the agent cannot be the monoclonal antibody alone, and pharmaceutically acceptable carrier.

A Alternative HER-2 transcript containing ECDIIIa sequence



B ECDIIIa HER-2 gene product:



p185 HER-2 gene product:




Figure 1

Figure 1 ; Doherty et al

Figure 2




Figure 3




Figure 4





Figure 5




Figure 6




Figure 7

DECLARATION AND POWER OF ATTORNEY

As the below-named inventor, we declare that we are the original, first, and sole inventors of the subject matter which is claimed in the specification identified below and for which a patent is sought on the invention as titled therein. We hereby state that we have reviewed the contents of said specification, including the claims. We acknowledge the duty to disclose information which is material to the examination of this application in accordance with 37 C.F.R. §1.56.

Inventor: Joni Kristin Doherty
 Address: 3928 SE Ogden Street
 Portland, Oregon 97202
 Residence: Portland, Oregon
 Citizenship: USA

Inventor: Gail M. Clinton
 Address: 3040 SW Beaverton Avenue
 Portland, Oregon 97201
 Residence: Portland, Oregon
 Citizenship: USA

Title of Invention: **HER-2 BINDING ANTAGONISTS**

Serial No.: _____, filed on January 20, 1999

(X) There are no earlier-filed United States Patent Applications of which priority benefit is claimed.

() We hereby claim the benefit under 35 U.S.C. §120 of the United States Patent Application listed below, and, insofar as the subject matter of the claims of this application is not disclosed in the prior United States application in the manner provided by the first paragraph of 35 U.S.C. §112, we acknowledge the duty to disclose material information as defined in 37 C.F.R. §1.56 which occurred between the filing date of the prior application and the filing date of this application:

USSN:	Filed:	Status:

POWER OF ATTORNEY

The power to prosecute this application and transact all business in the Patent and Trademark Office connected herewith is hereby granted to the following attorneys:

Jeffrey B. Oster, Registration No. 32,585
Todd M. Becker, Registration No. P43,487
Brian J. Deagle, Registration No. P43,579

Please send all correspondence to:

Jeffrey B. Oster
Davis Wright Tremaine
2600 Century Square
1501 Fourth Avenue
Seattle, Washington 98101-1688
Telephone: (206) 628-7711
Facsimile: (206) 628-7699

We further declare that all statements made herein of our own knowledge are true and that all statements made on information and belief are believed to be true; and further that these statements are made with the knowledge that willful false statements and the like so made are punishable by fine or imprisonment, or both, under Section 1001 of Title 18 of the United States Code, and that such willful false statements may jeopardize the validity of the application or any patent issuing thereon.

Signature of Inventor:

Date signed:

Joni Kristin Doherty

Gail M. Clinton